



**Rational Polypharmacy:  
An Update for Specific Conditions**

Thomas B. Gregory, PharmD, BCPS, CPE, FASPE

1

---

---

---

---

---

---

---

---

**Disclosures**

- Nothing to disclose



2

---

---

---

---

---

---

---

---

**In the news now...**

**Feds halt 2 Tennessee pharmacies' opioid dispensing for now**

By JONATHAN MATTISE February 8, 2019

The filings say Thomas Weir, who owns both pharmacies, oversaw operations and pharmacists.

Michael Griffith, John Dalton and I am not a doctor. I would like to see more doctors. Failure in the role. (a) A prescription for a controlled substance to be effective must be issued for a legitimate medical purpose by an individual practitioner acting in the usual course of his professional practice. The responsibility for the proper prescribing and dispensing of controlled substances is upon the prescribing practitioner, but a corresponding responsibility rests with the pharmacist who

fills the prescription. An order purporting to be a prescription issued not in the usual course of professional treatment or in legitimate and authorized research is not a prescription within the meaning and intent of section 309 of the Act (21 U.S.C. §29) and the person knowingly filling such a purported prescription, as well as the person issuing it, shall be subject to the penalties provided for violations of the provisions of law relating to controlled substances.

<https://apnews.com/fcae3106c7954369bf50905b6839ab6b>, accessed 3.6.2019  
[https://www.deadiversion.usdoj.gov/21cfr/cfr/1306/1306\\_04.htm](https://www.deadiversion.usdoj.gov/21cfr/cfr/1306/1306_04.htm), accessed 3.6.2019



3

---

---

---

---

---

---

---

---

**Learning Objectives**

- Define rational polypharmacy as it pertains to the patient in pain
- Recognize the various pharmacological classes used in rational polypharmacy of migraine, neuropathic pain, and musculoskeletal pain conditions
- Distinguish between rational and irrational polypharmacy in managing pain



4

---

---

---

---

---

---

---

---

**How does rational polypharmacy apply to my practice?**

- Synergistic combinations decreasing the amount of opioid needed for pain control
- Using nonopioids as first line therapy can minimize or even prevent the need for opioid medications on a chronic basis
- Shortages and regulatory constraints on the manufacture of opioids have lead to shortages and the inability of pharmacies to stock opioids and other medications used in pain management



5

---

---

---

---

---

---

---

---

**Definitions**

- Polypharmacy:  
The use of two or more drugs together, usually to treat a single condition or disease
- Synergy:  
The cooperative action of two or more stimuli or drugs
- Rational:  
Proceeding or derived from reason or based in reason
- Irrational:  
Not endowed with the faculty of reason



6

---

---

---

---

---

---

---

---

### Goals of Rational Polypharmacy

- Minimize adverse effects
  - Lower doses of individual medications
  - Opioid sparing effects
- Increase adherence to the prescribed regimen
- Using synergistic combinations of medications to achieve improved outcomes compared to the individual medications
- Increase efficacy by utilizing long acting and short acting preparations



7

---

---

---

---

---


---

---

---

### Hitting the Target(s)

- Stimulation of nociceptors causes signal transduction to the dorsal horn
  - Transduction
- The spinothalamic tract transmits the signals to the brain where pain is first experienced
  - Transmission and perception
- Descending pathways from the brain attempt to block the signal from the periphery
  - Modulation



8

---

---

---

---

---

---

---

---

### Medications Used in Pain Management

- Acetaminophen
- NSAIDs
- 5HT<sub>3</sub>-1B/D antagonists (Triptans)
- Calcitonin gene-related peptide antagonists
- Antidepressants
- Anticonvulsants
- Local anesthetics
- Skeletal muscle relaxants
- Opioids



9

---

---

---

---

---

---

---

---

### Acetaminophen

- Mechanism of action is still not entirely known
  - Thought to be a partial COX inhibitor
- March 2014 FDA mandates all prescription drug combination products containing acetaminophen cap the dose at 325 mg
- Maximum daily dose limits vary based on comorbidities and who you ask
  - FDA vs Johnson and Johnson

<http://www.fda.gov/drugs/drugsafety/informationbydrugclass/ucm165107.htm> accessed January 30, 2018  
<https://www.tylenol.com/safety-dosing/usage/dosage-for-adults> accessed January 30, 2018



10

---

---

---

---

---


---

---

---

### Acetaminophen (cont'd)

- Largest concern is unintentional overdoses
- Metabolism of acetaminophen by the liver is a saturable process
- Over the counter products and cumulative acetaminophen dosing



[http://www.medscape.com/viewarticle/518631\\_3](http://www.medscape.com/viewarticle/518631_3) accessed January 30, 2018



11

---

---

---

---

---

---

---

---

### Nonsteroidal Anti-Inflammatory Agents

- COX 1 more specific to the GI tract and renal homeostasis
- COX 2 more specific to inflammation and platelet aggregation
- Certain comorbidities limit the dosing on most NSAIDs
  - Patients on anticoagulants
  - Patients with renal dysfunction
  - Pregnancy



12

---

---

---


---

---

---

---

---



13

---

---

---

---

---


---

---

---

---

---



14

---

---

---

---

---

---

---

---

---

---

### Nonsteroidal Anti-Inflammatory Drugs

- Topical vs systemic NSAIDs
  - Patch, cream, lotion, etc
    - Range in application frequency from twice to four times daily
  - Topical can provide NSAID relief at the site of inflammation without the systemic side effects
  - Cost can be a limiting factor
  - Still carry a black box warning on the labeling for cardiovascular complications

15

---

---

---

---

---

---

---

---

---

---

### 5HT<sub>3-1B/D</sub> Antagonists (Triptans)

- Serotonin receptor antagonists leading to
  - Extra-cerebral vasoconstriction (5-HT<sub>1B</sub>)
  - Decreased inflammatory neuropeptide release (5-HT<sub>1D</sub>)
- Indicated for migraine treatment
  - Abortive therapy, not prophylactic
- Dosing in general involves administration of a second dose in 1 to 2 hours if the first dose was unsuccessful in aborting the migraine



16

---

---

---

---

---

---

---

---

### Triptans (cont'd)

Drug	Almotriptan	Eletriptan	Frovatriptan	Naratriptan	Rizatriptan	Sumatriptan	Zolmitriptan
Brand Name (Manufacturer)	Almotriptan (Lundbeck)	Relspan (Pfizer)	Frova (Janssen)	Amerge (SRI)	Maxalt, Maxalt MLT (Merck)	Imovio (UCB) Dorivac, Xolol (Ampel) Sumatriptan (Eli Lilly) Zembrace Sym-Touch (Pharmacia)	Zemig, Zemig Start (Drugs)
Generic Available	Yes	No	Yes	Yes	Yes	Yes – for generic products only	Yes – for oral tabs and ODTs only
Route of Administration	Oral	Oral	Oral	Oral	Oral	Oral, Nasal, SQ	Oral, Nasal
Formulations	6, 25, 50, 100 mg tabs	25, 40 mg tabs	2.5 mg tabs	1, 2.5 mg tabs	5, 10 mg tabs and 5, 10 mg ODTs	Injection and generic – Oral 25, 50, 100 mg tabs; SQ 4, 8 mg/0.5 mL auto-injector pen and self-injector auto; Nasal 2, 5 mg/0.5 mL nasal spray; Dorivac Xolol 11 mg nasal powder sacs; Sumatriptan 4 mg/0.5 mL SQ; Zolmitriptan 2.5 mg/0.5 mL SQ auto-injector	Oral 2.5 mg tabs and 2.5 mg ODTs; Nasal 2.5 mg/0.5 mL nasal spray
Onset of Action	30-60 min	30-60 min	~3 hrs	1-3 hrs	30-60 min	Take 30-60 min SQ ~10 min; Nasal 30-15 min	Take 30-60 min; Nasal 10-15 min
Elimination Half-life	1.5-4 hrs	~4 hrs	~25 hrs	~6 hrs	2-3 hrs	~2 hrs	2-3 hrs

<http://www.headache.mobi/uploads/1/1/7/5/11757140/triptans.pdf> accessed 2.28.2019



17

---

---

---

---

---

---

---

---

### Triptans (cont'd)

- Patients that are NOT candidates for triptan agents
  - Ischemic heart disease
  - Uncontrolled hypertension
  - Peripheral vascular disease
  - History of cerebrovascular syndromes (stroke or transient ischemic attack)
- Multiple formulations exist for
  - Sumatriptan (nasal, SQ, oral)
  - Zolmitriptan (nasal and oral)



18

---

---

---

---

---

---

---

---

### Calcitonin Gene-Related Peptide (CGRP) Antagonists

- Monoclonal antibodies that bind to CGRP
  - Preventing intracranial artery vasodilatation
  - Prevention of dural mast cell degranulation
- Indicated for the prevention of migraine
  - Not indicated for the management of acute migraine symptoms
- Administration of the currently approved agents monthly subcutaneous injection



AnnRevPharmacolTox.55.533-52 2015

19

---

---

---

---

---

---

---

---

### CGRP Antagonists Currently Available

- Erenumab-aooe [Aimovig®]
  - Subcutaneous injection 70 mg once monthly
  - May increase to 70 mg twice a month in some patients
- Fremanezumab-vfrm [Ajovy®]
  - Subcutaneous injection 225 mg once monthly or 675 mg every three months
- Galcanezumab-gnlm [Emgality®]
  - Subcutaneous injection 240 mg once then 120 mg monthly



Lexicomp accessed 3.1.2019

20

---

---

---

---

---


---

---

---

### CGRP Antagonists (cont'd)

- Questions that remain unanswered regarding their long term safety include
  - Hypertension
  - Nitric oxide synthase
  - Platelet aggregation
  - Negative impact on microvasculature
    - Heart failure
    - Diabetes



<https://www.practicalpainmanagement.com/pain/headache/state-possible-long-term-side-effects-cgrp-antagonists>, accessed 3.1.2019

21

---

---

---

---

---

---

---

---

**Tricyclic Antidepressants (TCA)**

- Mechanism of action is through inhibition of norepinephrine and serotonin reuptake and inhibition of sodium channel action potentials
- The antidepressant effects and the neuropathic pain analgesia are independent
  - Higher dosing and longer treatment time needed for antidepressant effects
- Caution should be exercised in patients
  - With cardiac arrhythmias
  - Over the age of 65



22

---

---

---

---

---

---

---

---

**Serotonin Norepinephrine Reuptake Inhibitors (SNRI)**

- Mechanism of action is through inhibition of norepinephrine and serotonin reuptake
- Dosing is generally higher for treating neuropathic pain compared to treating depression
- Withdrawal syndromes can occur if patients are taken off SNRI therapy abruptly
  - Anxiety, irritability, headache, paresthesia, nervousness
- Caution should be exercised in patients with liver dysfunction, uncontrolled hypertension, or moderate cardiovascular disease



23

---

---

---

---

---

---

---

---

**Antiepileptics**

- The primary antiepileptics used in pain management work on calcium channels
  - Gabapentin
  - Pregabalin
- Other antiepileptics have had mixed results regarding neuropathic pain
  - Valproic acid
  - Phenytoin
- Carbamazepine for trigeminal neuralgia



24

---

---

---

---

---

---

---

---



**Local Anesthetics**

- Mechanism of action is through membrane stabilization of sodium channels preventing depolarization and signal transduction
- Acute uses for local anesthesia (procedures, etc)
  - Topical application
    - Cream, ointment, patch, etc
  - Intradermal injections
  - Nerve blocks
- Patches are indicated for the management of postherpetic neuralgia



25

---

---

---

---

---

---

---

---

**Skeletal Muscle Relaxants**

- Multiple medications are included in this general taxonomy
  - Certain agents approved for spasticity
    - Baclofen and tizanidine
- Others stand out for reasons other than their indication
  - Cyclobenzaprine and orphenadrine regarding their anticholinergic effects
  - Chlorzoxazone and potential for hepatotoxicity
  - Carisopradol and meprobamate and potential for abuse



26

---

---

---

---

---

---

---

---

**Opioids**

- Opioids work on multiple receptors within the CNS
  - Analgesia and adverse effects are derived from mostly mu receptors
- There is no ceiling dose for analgesia; however, as doses increase the incidence of adverse effects increases
- CDC (2016) and VA/DoD (2017) guidelines outlining the use of opioids in chronic pain have been published



27

---

---

---

---

---

---

---

---

**Opioids (cont'd)**

- Agonists vs partial agonists vs antagonists
  - Morphine, fentanyl, methadone, etc
  - Buprenorphine, nalbuphine, butorphanol
  - Naloxone and naltrexone
  
- Awareness of other nonpain combination products
  - Naltrexone-bupropion for weight loss



28

---

---

---

---

---


---

---

---

**Opioid Statistics**

- Medication overdose deaths in 2016: 63,632
  - Opioids (illicit and prescription) were involved in 66.4% of those fatalities
  
- Patients on > 90 morphine milligram equivalents have decreased from 11.5 to 5 per 100 patients in the US



<https://www.cdc.gov/drugoverdose/pdf/2016-cdc-drug-surveillance-report.pdf#page=72> accessed 3.6.2019



29

---

---

---

---


---

---

---

---

**Opioid Statistics (cont'd)**



<https://www.cdc.gov/drugoverdose/pdf/2016-cdc-drug-surveillance-report.pdf#page=75> accessed 3.6.2019



30

---

---

---


---

---

---

---

---



31

---

---

---

---

---

---

---

---

---

---

- ### Patients at Risk for Opioid Adverse Events
- Patients with sleep apnea and sleep disordered breathing
  - Pregnancy
  - Hepatic or renal dysfunction
  - Age greater than 65
  - Mental health or substance use disorders
  - Nonfatal overdose history

32

---

---

---

---

---


---

---

---

---

---



33

---

---

---

---

---

---

---

---

---

---

**Immediate Release (IR) vs Extended Release (ER)**

- Initial therapy should include the use of IR formulations
- ER preparations are appropriate for patients
  1. That routinely use the IR preparation with relief of pain
  2. That are not experiencing adverse effects that decrease quality of life
  3. That are on stable doses of IR preparations and have been for an appropriate time frame
- IR and ER preparation use should be re-evaluated for safety and efficacy periodically or per state guideline



34

---

---

---

---

---

---

---

---

**Nonrational Polypharmacy**

- Utilizing two medications in the same family for the same condition
  - Ibuprofen and naproxen
  - Morphine immediate release and oxycodone immediate release
- Adding a medication that may be contraindicated based on the patients other comorbidities
  - Methadone use in a patient with a history of QTc prolongation
  - Tramadol or use in a patient with underlying seizure history



35

---

---

---

---

---

---

---

---

**Rationalizing Migraine Pain Management**

- Use of abortive medications at the beginning of a migraine
  - NSAIDs, triptans
  - Opioids and dopamine antagonists (severe)
- Use of prophylactic therapy once patients meet criteria
  - More than two migraines per month
  - Migraine lasts for more then 24 hours
  - Use of abortive therapy more than twice per week



36

---

---

---

---

---

---

---

---

### Rationalizing Migraine PROPHYLACTIC Management

		Common Condition	Medication
First line	High efficacy	Beta blockers	
		Tricyclic antidepressants	
	Low efficacy	Verapamil	
Second line	High efficacy	Metylergoline	
		Flunarizine	
	Unproven efficacy	MAOIs	
		Cyclophosphamide	
		Depression	Tricyclic antidepressants, SSRIs
		Overweight	Topiramate, propranolol
		Underweight	Tricyclic antidepressants (nortriptyline, protriptyline)
		Epilepsy	Valproic acid, topiramate
		Mastic	Valproic acid

MAOIs = monoamine oxidase inhibitors  
SSRIs = selective serotonin reuptake inhibitors

**PainWeek** <https://medicine.medscape.com/article/1142556-treatment> accessed 3.4.2019

37

---

---

---

---

---

---

---

---

### Rationalizing Neuropathic Pain

- Scheduled use of tricyclic or SNRI antidepressants at appropriate doses
  - Caution regarding the use of anticholinergic tricyclic agents
- Use of antiepileptics at appropriate doses
  - Opioids may be used in combination with the use of an antiepileptic
  - Topical local anesthetics such as patches and creams with the above

**PainWeek**

38

---

---

---

---

---

---

---

---

### Rationalizing Neuropathic Pain (cont'd)

- NSAIDs and acetaminophen are unlikely to alleviate neuropathic pain
- Anticonvulsants, local anesthetics, and tricyclic antidepressants are mainstays in neuropathic pain management
- Opioids may have a place but not first or second line
- Muscle relaxants are controversial in terms of efficacy

Medication	Dose	Notes	Side Effects
Tricyclic antidepressants (TCAs)	75-150 mg/day	Start with 75 mg at bedtime and increase by 25 mg every 1-2 weeks. Do not exceed 150 mg/day. Avoid in patients with urinary retention, glaucoma, or severe constipation.	Anticholinergic effects, sedation, weight gain, orthostatic hypotension, cardiac conduction system effects.
Serotonin-norepinephrine reuptake inhibitors (SNRIs)	75-150 mg/day	Start with 75 mg at bedtime and increase by 25 mg every 1-2 weeks. Do not exceed 150 mg/day.	Serotonin toxicity, hypertension, dry mouth, constipation.
Anticonvulsants	Varies by drug	Carbamazepine: 800-1200 mg/day. Gabapentin: 3000-6000 mg/day. Pregabalin: 300-600 mg/day. Lamotrigine: 200-300 mg/day. Topiramate: 600-900 mg/day.	Dizziness, ataxia, blurred vision, hyponatremia, weight loss, hair loss.
Local anesthetics	Varies by drug	Lidocaine patches: 5% lidocaine patches. Capsaicin patches: 8% capsaicin patches.	Local irritation, systemic toxicity (rare).
Opioids	Varies by drug	Start with low doses and titrate up. Avoid in patients with respiratory depression or severe constipation.	Respiratory depression, constipation, dependence.
Muscle relaxants	Varies by drug	Cyclobenzaprine: 30 mg/day. Baclofen: 15-30 mg/day. Tizanidine: 4-8 mg/day.	Sedation, dizziness, hypotension, dry mouth.

**PainWeek** <https://www.uspharmacist.com/article/postherpetic-neuralgia-seniors-at-risk> accessed 3.5.2019

39

---

---

---

---

---


---

---

---

### Rationalizing Musculoskeletal Pain Management

- Bone pain
  - Muscle pain
  - Tendon and ligament pain
  - Fibromyalgia
  - Joint pain
  - Nerve compression syndromes
- More than 150 diagnoses all of which affect the locomotor system



<https://pmj.bmj.com/content/79/937/627>  
accessed 3.7.2019



40

---

---

---

---

---

---

---

---

---

---

### Rationalizing Musculoskeletal Pain Management (cont'd)

	Key OA
Strongly Recommended	None
Conditionally Recommended	<ul style="list-style-type: none"> <li>• Acetaminophen</li> <li>• NSAIDs</li> <li>• Tramadol</li> <li>• Non-narcotic combination analgesics</li> </ul>
Conditionally Not Recommended	<ul style="list-style-type: none"> <li>• Opioids</li> <li>• Gabapentin</li> </ul>
No Recommendation	<ul style="list-style-type: none"> <li>• Spinal NSAIDs</li> <li>• Intra-articular hyaluronic acid injections</li> <li>• Corticosteroids</li> <li>• Spinal analgesics</li> </ul>
	Rheu OA
Strongly Recommended	None
Conditionally Recommended	<ul style="list-style-type: none"> <li>• Acetaminophen</li> <li>• NSAIDs</li> <li>• Tramadol</li> <li>• Non-narcotic combination analgesics</li> </ul>
Conditionally Not Recommended	<ul style="list-style-type: none"> <li>• Opioids</li> <li>• Gabapentin</li> </ul>
No Recommendation	<ul style="list-style-type: none"> <li>• Spinal NSAIDs</li> <li>• Intra-articular hyaluronic acid injections</li> <li>• Corticosteroids</li> <li>• Spinal analgesics</li> </ul>

ACR: American College of Rheumatology; NSAID, non-steroidal anti-inflammatory drug; OA, osteoarthritis.

<https://www.practicalpainmanagement.com/treatment-osteoarthritis>, accessed 3.7.2019



41

---

---

---

---

---

---

---

---

---

---

### Rationalizing Musculoskeletal Pain Management (cont'd)

Drug (Brand)	Class	Status	Dosing
Amitriptyline (Elavil)	TCA	OTC status	Initiate at 10 mg po qhs, may titrate up to 50 mg, 2 months
Duloxetine (Cymbalta)	SNRI	FDA-approved	Initiate at 30 mg po qd for 1 wk, then increase to 60 mg qd
Milnacipran (Savella)	SNRI	OTC status	Initiate at 10 mg po qhs, then slowly titrate to 10 mg po bid and 20 mg po bid
Pregabalin (Lyrica)	GABA agonist	OTC status	Initiate at 30 mg po bid and gradually titrate up to response, max 300 mg qd
Gabapentin (Neurontin)	GABA agonist	OTC status	Initiate at 15 mg po bid, may increase to 150 mg bid based on tolerability and response, max 300 mg bid
Tramadol (Ultram)	SNRI	OTC status	Initiate at 300 mg po over 1 wk, after 1 wk, gradually increase to target dose of 400 mg qd, titrate according to response
Milnacipran (Savella)	SNRI	OTC status	Initiate at 10 mg po bid, may increase to 150 mg bid based on tolerability and response, max 300 mg bid
Venlafaxine (Effexor XR)	SNRI	OTC status	Initiate at 37.5 mg po qd or 75 mg po bid, may increase to 150 mg po qd or 300 mg po bid
Tramadol (Ultram)	SNRI	OTC status	Initiate at 30 mg po qd, then titrate to 25 mg increments on response, avoid level 2 drugs to reach 100 mg/day, go to double-blind (25 mg po qd), after titration, 50-100 mg po qd po qd, up to 400 mg/day po qd

OTC: over-the-counter; TCA, tricyclic antidepressant; SNRI, serotonin-norepinephrine reuptake inhibitor; SNRI, selective norepinephrine reuptake inhibitor; TCA, tricyclic antidepressant; TCA, tricyclic antidepressant.

<https://www.uspharmacist.com/article/treatment-of-fibromyalgia-pain>, accessed 3.7.2019



42

---

---

---

---

---

---

---

---

---

---

**Conclusion**

- Pain management typically involves more than one modality in order to manage
- Safety must take into consideration patient specific factors that will change over time
- Certain combinations can put patients at risk for adverse effects but having a complete picture of a patients medications can help prevent this



43

---

---

---

---

---

---

---

**See you at PAINWEEK**



44

---

---

---

---

---

---

---