



Rational Polypharmacy: An Update for Specific Conditions

Thomas B. Gregory, PharmD, BCPS, CPE, FASPE

Disclosures

- Nothing to disclose



In the news now...

Feds halt 2 Tennessee pharmacies' opioid dispensing for now

By JONATHAN MATTISE February 8, 2019

The filings say Thomas Weir, who owns both pharmacies, oversaw operations and pharmacists

Michael Griffith, John Polston and Laura Larkin Health filed reservations, falling in the role

(e) A prescription for a controlled substance to be effective must be issued for a legitimate medical purpose by an individual practitioner acting in the usual course of his professional practice. The responsibility for the proper prescribing and dispensing of controlled substances is upon the prescribing practitioner, but a corresponding responsibility rests with the pharmacist who dispenses the substance. An order purporting to be a prescription issued not in the usual course of professional treatment or in legitimate and authorized research is not a prescription within the meaning and intent of section 309 of the Act (21 U.S.C. 829) and the person knowingly filling such a purported prescription, as well as the person issuing it, shall be subject to the penalties provided for violations of the provisions of law relating to controlled substances.

<https://apnews.com/fcae3106c7954369bf50905b6639ab6b> accessed 3.6.2019
https://www.deadiversion.usdoj.gov/21cfr/cfr/1306/1306_04.htm accessed 3.6.2019



Learning Objectives

- Define rational polypharmacy as it pertains to the patient in pain
- Recognize the various pharmacological classes used in rational polypharmacy of migraine, neuropathic pain, and musculoskeletal pain conditions
- Distinguish between rational and irrational polypharmacy in managing pain



How does rational polypharmacy apply to my practice?

- Synergistic combinations decreasing the amount of opioid needed for pain control
- Using nonopioids as first line therapy can minimize or even prevent the need for opioid medications on a chronic basis
- Shortages and regulatory constraints on the manufacture of opioids have lead to shortages and the inability of pharmacies to stock opioids and other medications used in pain management



Definitions

- Polypharmacy:
The use of two or more drugs together, usually to treat a single condition or disease
- Synergy:
The cooperative action of two or more stimuli or drugs
- Rational:
Proceeding or derived from reason or based in reason
- Irrational:
Not endowed with the faculty of reason




Goals of Rational Polypharmacy

- Minimize adverse effects
 - Lower doses of individual medications
 - Opioid sparing effects
- Increase adherence to the prescribed regimen
- Using synergistic combinations of medications to achieve improved outcomes compared to the individual medications
- Increase efficacy by utilizing long acting and short acting preparations



Hitting the Target(s)

- Stimulation of nociceptors causes signal transduction to the dorsal horn
 - Transduction
- The spinothalamic tract transmits the signals to the brain where pain is first experienced
 - Transmission and perception
- Descending pathways from the brain attempt to block the signal from the periphery
 - Modulation



Medications Used in Pain Management

- Acetaminophen
- NSAIDs
- 5HT_{3-1B/D} antagonists (Triptans)
- Calcitonin gene-related peptide antagonists
- Antidepressants
- Anticonvulsants
- Local anesthetics
- Skeletal muscle relaxants
- Opioids



Acetaminophen


- Mechanism of action is still not entirely known
 - Thought to be a partial COX inhibitor
- March 2014 FDA mandates all prescription drug combination products containing acetaminophen cap the dose at 325 mg
- Maximum daily dose limits vary based on comorbidities and who you ask
 - FDA vs Johnson and Johnson

<http://www.fda.gov/drugs/drugsafety/informationbydrugclass/ucm165107.htm> accessed January 30, 2018
<https://www.tylenol.com/safety-dosing/usage/dosage-for-adults> accessed January 30, 2018



Acetaminophen (cont'd)

- Largest concern is unintentional overdoses
- Metabolism of acetaminophen by the liver is a saturable process
- Over the counter products and cumulative acetaminophen dosing




http://www.medscape.com/viewarticle/518631_3 accessed January 30, 2018




Nonsteroidal Anti-Inflammatory Agents

- COX 1 more specific to the GI tract and renal homeostasis
- COX 2 more specific to inflammation and platelet aggregation
- Certain comorbidities limit the dosing on most NSAIDs
 - Patients on anticoagulants
 - Patients with renal dysfunction
 - Pregnancy







Nonsteroidal Anti-Inflammatory Drugs

- Topical vs systemic NSAIDs
 - Patch, cream, lotion, etc
 - Range in application frequency from twice to four times daily
 - Topical can provide NSAID relief at the site of inflammation without the systemic side effects
 - Cost can be a limiting factor
 - Still carry a black box warning on the labeling for cardiovascular complications

Calcitonin Gene-Related Peptide (CGRP) Antagonists

- Monoclonal antibodies that bind to CGRP
 - Preventing intracranial artery vasodilatation
 - Prevention of dural mast cell degranulation
- Indicated for the prevention of migraine
 - Not indicated for the management of acute migraine symptoms
- Administration of the currently approved agents monthly subcutaneous injection



AnnRevPharmacolTox.55.533-52.2015

CGRP Antagonists Currently Available


- Erenumab-aooe [Aimovig®]
 - Subcutaneous injection 70 mg once monthly
 - May increase to 70 mg twice a month in some patients
- Fremanezumab-vfrm [Ajovy®]
 - Subcutaneous injection 225 mg once monthly or 675 mg every three months
- Galcanezumab-gnlm [Emgality®]
 - Subcutaneous injection 240 mg once then 120 mg monthly



Lexicomp accessed 3.1.2019

CGRP Antagonists (cont'd)

- Questions that remain unanswered regarding their long term safety include
 - Hypertension
 - Nitric oxide synthase
 - Platelet aggregation
 - Negative impact on microvasculature
 - Heart failure
 - Diabetes



<https://www.practicalpainmanagement.com/pain/headache/slake-possible-long-term-side-effects-cgrp-antagonists> accessed 3.1.2019



Tricyclic Antidepressants (TCA)

- Mechanism of action is through inhibition of norepinephrine and serotonin reuptake and inhibition of sodium channel action potentials
- The antidepressant effects and the neuropathic pain analgesia are independent
 - Higher dosing and longer treatment time needed for antidepressant effects
- Caution should be exercised in patients
 - With cardiac arrhythmias
 - Over the age of 65



Serotonin Norepinephrine Reuptake Inhibitors (SNRI)

- Mechanism of action is through inhibition of norepinephrine and serotonin reuptake
- Dosing is generally higher for treating neuropathic pain compared to treating depression
- Withdrawal syndromes can occur if patients are taken off SNRI therapy abruptly
 - Anxiety, irritability, headache, paresthesia, nervousness
- Caution should be exercised in patients with liver dysfunction, uncontrolled hypertension, or moderate cardiovascular disease



Antiepileptics

- The primary antiepileptics used in pain management work on calcium channels
 - Gabapentin
 - Pregabalin
- Other antiepileptics have had mixed results regarding neuropathic pain
 - Valproic acid
 - Phenytoin
- Carbamazepine for trigeminal neuralgia



Local Anesthetics

- Mechanism of action is through membrane stabilization of sodium channels preventing depolarization and signal transduction
- Acute uses for local anesthesia (procedures, etc)
 - Topical application
 - Cream, ointment, patch, etc
 - Intradermal injections
 - Nerve blocks
- Patches are indicated for the management of postherpetic neuralgia



Skeletal Muscle Relaxants

- Multiple medications are included in this general taxonomy
 - Certain agents approved for spasticity
 - Baclofen and tizanidine
- Others stand out for reasons other than their indication
 - Cyclobenzaprine and orphenadrine regarding their anticholinergic effects
 - Chlorzoxazone and potential for hepatotoxicity
 - Carisoprodol and meprobamate and potential for abuse



Opioids

- Opioids work on multiple receptors within the CNS
 - Analgesia and adverse effects are derived from mostly mu receptors
- There is no ceiling dose for analgesia; however, as doses increase the incidence of adverse effects increases
- CDC (2016) and VA/DoD (2017) guidelines outlining the use of opioids in chronic pain have been published



Opioids (cont'd)

- Agonists vs partial agonists vs antagonists
 - Morphine, fentanyl, methadone, etc
 - Buprenorphine, nalbuphine, butorphanol
 - Naloxone and naltrexone


- Awareness of other nonpain combination products
 - Naltrexone-bupropion for weight loss



Opioid Statistics

- Medication overdose deaths in 2016: 63,632
 - Opioids (illicit and prescription) were involved in 66.4% of those fatalities


- Patients on > 90 morphine milligram equivalents have decreased from 11.5 to 5 per 100 patients in the US



https://www.cdc.gov/drugoverdose/pdf/nbs/2018_cdc_drug_surveillance_report.pdf#page=72 accessed 3.6.2019



Opioid Statistics (cont'd)



https://www.cdc.gov/drugoverdose/pdf/nbs/2018_cdc_drug_surveillance_report.pdf#page=75 accessed 3.6.2019



Immediate Release (IR) vs Extended Release (ER)

- Initial therapy should include the use of IR formulations
- ER preparations are appropriate for patients
 1. That routinely use the IR preparation with relief of pain
 2. That are not experiencing adverse effects that decrease quality of life
 3. That are on stable doses of IR preparations and have been for an appropriate time frame
- IR and ER preparation use should be re-evaluated for safety and efficacy periodically or per state guideline



Nonrational Polypharmacy

- Utilizing two medications in the same family for the same condition
 - Ibuprofen and naproxen
 - Morphine immediate release and oxycodone immediate release
- Adding a medication that may be contraindicated based on the patients other comorbidities
 - Methadone use in a patient with a history of QTc prolongation
 - Tramadol or use in a patient with underlying seizure history



Rationalizing Migraine Pain Management

- Use of abortive medications at the beginning of a migraine
 - NSAIDs, triptans
 - Opioids and dopamine antagonists (severe)
- Use of prophylactic therapy once patients meet criteria
 - More than two migraines per month
 - Migraine lasts for more than 24 hours
 - Use of abortive therapy more than twice per week



Conclusion

- Pain management typically involves more than one modality in order to manage
- Safety must take into consideration patient specific factors that will change over time
- Certain combinations can put patients at risk for adverse effects but having a complete picture of a patients medications can help prevent this



See you at PAINWEEK