



Testing the Waters: Urine Drug Screening for the Perplexed Among Us

Mark Garofoli, PharmD, MBA, BCGP, CPE

1

Faculty



- Family of 8 Pharmacists
- Wife, In-Laws, & Cousins
- Family Vineyard in the Marche Region of Italy



2

Disclosures

- Nothing to disclose

This presentation was not a part of the presenter's official duties at the WVU and does not represent the opinion of WVU



3

Learning Objectives

- Identify the clinical practice implication differences between urine drug screenings and urine drug tests
- Describe anticipated results of urine drug screenings and urine drug tests, based on prescribed medications and utilized substances
- Recall common cross reactants for controlled substances in urine drug screenings




4

2016 CDC Chronic Pain Opioid Guidelines

5

2016 CDC Chronic Pain Opioid Guidelines



6

TRUST, but Verify



Painweek

7

Urine Drug Monitoring (UDM)

Urine Drug Screening (UDS)

Urine Drug Testing (UDT)

Painweek

8

Urine Drug Monitoring



Goals

- Improve proper medication adherence
- Prevent ± detect medication misuse/diversion

Painweek

Wheeler, K. et al. © Urine Drug Screening: Practical Guide for Clinicians, Mayo Clin Proc. 2008;83(9):1171-7

9

Urine Drug Monitoring: Frequency

Risk	Frequency
Low	Annual
Moderate	≥2x/Year
High	≥3x/Year
Any	? Every Appointment ?



Adapted from: Mueller, K. et al. Urine Drug Screening: Practical Guide for Clinicians. Mayo Clin Proc. 2008;83(4):672-76

10

Urine Drug Screening Strategies

- Patient/provider agreement → UDM procedures
- Randomized or scheduled (eg, appointments)
- Urine samples collected in a private bathroom without running water, soap, hand sanitizer, or other liquids, and with toilet water stained blue
- Urine specimen cups with temperature strips that fluoresce between 90°F to 100°F
- Urine creatinine and specific gravity can be ordered together with a drug test panel



Mueller, K. et al. Urine Drug Screening: Practical Guide for Clinicians. Mayo Clin Proc. 2008;83(4):672-76

11

Debunking UDS Mythical Work-Arounds

- 1. Home remedies**
–Niacin, bleach, certo, vinegar, goldenseal, cranberry juice
- 2. Synthetic urine**
- 3. Dilution**
- 4. Substitution**
–Someone else's urine



Mueller, K. et al. Urine Drug Screening: Practical Guide for Clinicians. Mayo Clin Proc. 2008;83(4):672-76

12

Online Urine Purchasing
www.thewhizzinator.com

The Whizzinator Touch in White
 The Whizzinator Touch in Tan
 The Whizzinator Touch in Latine
 The Whizzinator Touch in Brown
 The Whizzinator Touch in Black

Painweek

13

Homemade Urine Recipe(s)

Recipe 1

- 976 ml of 0.02 mole hydrochloric acid solution
- 1.9 ml of 0.25 ammonia solution
- 14.1 g sodium chloride or table salt
- 2.8 g potassium chloride
- 17.3 g urea
- 0.6 g calcium chloride
- 0.43 g magnesium sulfate

Recipe 2

- 750 ml distilled water
- 7.5 g sodium chloride or table salt
- 50 mg albumin powder (bovine or egg)
- 4.5 g potassium chloride
- 4.8 g sodium phosphate (monobasic)
- 18.2 g urea
- 2 g creatinine

Painweek

14

Urine Color

- ❖ The yellow coloration of urine results from urobilin that is produced as a product of bilirubin degradation
- ❖ Normal urine color: light yellow to golden

AM I HYDRATED?
Urine Color Chart

1		If your urine matches this color, you are drinking enough fluids.
2		Drink more water to get the ideal color to shade 1 and 3.
3		Dehydrated
4		You may suffer from cramps and heat-related problems.
5		Health risk! Drink more water.
6		Health risk! Drink more water.
7		Health risk! Drink more water.
8		Health risk! Drink more water.

Painweek

15

Urine Color	Medications	Description
Orange	Chlorzoxazone, isoniazid, phenazopyridine, sulfasalazine, & warfarin	Consumption of carrots Monitor for sparse blood in urine (hematuria)
Red	Chlorzoxazone, ibuprofen, phenazopyridine, rifampin, senna, & warfarin	Consumption of red beets, rhubarb, or carrots Monitor for sparse blood in urine (hematuria) Myoglobinuria from rhabdomyolysis
Brown	Acetaminophen, metronidazole, & nitrofurantoin	Myoglobinuria from rhabdomyolysis ("hand drumming") Acute renal/hepatic disease Metastatic melanoma (rare reports)
Purple	Chlorzoxazone Combination of medications causing red or blue	Gram-negative bacteria
Blue	Amitriptyline, cimetidine, indomethacin, methocarbamol, metoclopramide, & zaleplon	Methylene blue
Green	Methocarbamol Medications causing blue (added to yellow urine)	Consumption of asparagus or black licorice UTI with pseudomonas
White	x	Calcium or phosphate crystals infection
Black	Methocarbamol, methyldopa/L-dopa, senna, & sorbitol	Phenol or copper poisoning Consumption of iodine Metastatic melanoma (rare reports)

Painweek

08. 8. et al. JAMA Dermatol. 2014; 150(8):1273-1274

16

Biologic Specimen Detection

Biologic Specimen	Detection Time After Ingestion					
	Minutes	Hours	Days	Weeks	Months	Years
Blood	X	X				
Saliva	X	X				
Urine	X	X	X			
Sweat	X	X	X	X		
Hair			X	X	X	X
Nails			X	X	X	X

Painweek

Caplan YH et al. J Anal Toxicology 2011; 25: 296-299

17

Urine Drug Detection Times

Drug	Detection Time After Ingestion
Alcohol	7 to 12 hours
Amphetamines	2 to 3 days
Benzodiazepines (short-acting)	3 days
Benzodiazepines (long-acting)	30 days
Marijuana (single dose)	3 days
Marijuana (4x/week)	5 to 7 days
Marijuana (daily)	10 to 15 days
Marijuana (long-term)	>30 days
Codeine	2 days
Heroin	2 days
Hydromorphone	2 to 4 days
Methadone	3 days
Morphine	2 to 3 days
Oxycodone	2 to 4 days

Painweek

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3400000/pdf/nihms248887.pdf


18

URINE DRUG MONITORING	
Urine Drug Screening (UDS)	Urine Drug Testing (UDT)
Immunoassay screen (ie, cup)	GC-MS or LC-MS
PRESUMPTIVE	DEFINITIVE
In-office, point-of-care, or lab-based	Laboratory, highly specific & sensitive
Results within minutes	Results in hours or days
Various cups detect a majority of legal & illicit medications by structural class	Measures all drug/metabolite concentrations
Guidance for preliminary treatment decisions	Definitive identification & analysis
Cross reactivity common: more false positives	False-positive results are rare
Higher cutoff levels: more false negatives	False-negative results are rare
\$	\$\$\$

Painweek

Wheeler, K. et al. Urine Drug Screening: Practical Guide for Clinicians. West Co. Press, 2008. 101-107.

19



20

One Fancy Cup Immunoassay

INDIVIDUAL TESTS

Test Dipped In Urine

READ RESULTS

C POSITIVE, [] NEGATIVE, [] NEGATIVE, [] NEGATIVE

Painweek

21

Urine Drug Monitoring

Cut-Off Amounts (SAMSHA)

Chemical	UDS Cut-Off (ng/mL)	UDT Cut-Off (ng/mL)
THC	50	15
Opiates	2,000	2,000
Hydrocodone/hydromorphone	300	100
Oxycodone/oxymorphone	100	100
6-MAM	10	10
Amphetamines/meth/MDMA	500	250
Cocaine (benzoylecgonine)	150	100
PCP	25	25

Painweek © 2019 Painweek, LLC. All rights reserved. For personal use only. For more information, please contact us at 800-877-6789 or www.painweek.com

22

Opioid Structural Classes

Phenanthrenes		Benzomorphans	Phenylpiperidines	Dipheylheptanes	Phenylpropylamines
5 Rings	4 Rings	3 Rings	2 Rings	2 Rings	2 Rings
Buprenorphine Codeine Diacetylmorphine Hydrocodone Morphine Naloxone Oxycodone Oxymorphone	Butorphanol Levorphanol	Diphenoxylate Loperamide Pentazocine	Fentanyl Meperidine	Methadone Propoxyphene	Tapentadol Tramadol

Painweek Adapted from Yalkow MD, Yalkow AT. NMP1 2014. EN1283-1283

23

URINE DRUG SCREENS

SUBSTANCE	5 Panel	7 Panel	10 Panel	12 Panel	13 Panel	14 Panel
THC	X	X	X	X	X	X
Cocaine	X	X	X	X	X	X
Opiates	X	X	X	X	X	X
PCP	X	X	X	X	X	X
Amphetamines	X	X	X	X	X	X
Benzodiazepines		X	X	X	X	X
Barbiturates		X	X	X	X	X
Methadone			X	X	X	X
Propoxyphene			X	X	X	X
Quaaludes			X	X	X	X
Ecstasy				X	X	X
Oxycodone				X	X	X
Fentanyl					X	X
Meperidine					X	X
Buprenorphine						X

Painweek Stallin, K. et al. Urine Drug Screenings: Practical Guide for Clinicians. Mayo Clin Proc. 2008;83:1595-1610. <http://www.mayoclinic.com/association/urine-drug-screening-panel-for-2008-08-08>


24

Urine Drug Screenings (UDS) Add-On Screenings

6-MAM (6-AM)
Synthetic cannabinoids
Cathinones (bath salts)
Dextromethorphan (DXM)
Ethanol
Propofol
Methylphenidate
Carisoprodol
Zolpidem
And so on...

PainWeek

25




26

Urine Drug Screening False Positives	
Substance	UDS Cross-Reactant
Cannabinoids	Dronabinol, NSAIDs (ibuprofen/naproxen), efavirenz, PPIs (pantoprazole), & promethazine
Opioids	chlorpromazine, dextromethorphan, diphenhydramine, doxylamine, poppy seeds, quinine, & quinolones, rifampin, & verapamil
Amphetamines	amantadine, bupropion, chlorpromazine, fluoxetine, labelalol, menthol, metformin, methylphenidate, phentermine, phenylephrine, promethazine, propranolol, pseudoephedrine, ranitidine, selegiline (L-meth/amphetamine metabolites), & trazodone
PCP	chlorpromazine, dextromethorphan, diphenhydramine, doxylamine, ibuprofen, imipramine, lamotrigine, meperidine, thioridazine, tramadol, & venlafaxine
LSD	amitriptyline, bupropion, buspirone, diltiazem, fentanyl, fluoxetine, labelalol, methylphenidate, metoprolamide, prochlorperazine, risperidone, sertraline, trazodone, & verapamil
Barbiturates	ibuprofen & naproxen
Benzodiazepines	oxaprozin, sertraline, & some herbals
Alcohol	asthma inhalers & isopropyl alcohol (short chain alcohol)
Methadone	quetiapine
TCAs	carbamazepine, cyclobenzaprine, diphenhydramine, hydroxyzine, & quetiapine


PainWeek

27

Urine Drug Monitoring



Conversation Starters



Conversation Leaders

Painweek

28

UDM Billing Terminology

- **ELISA or EIA**
 - Enzyme-linked immunosorbent assay
 - Detects and measures antibodies in your blood
- **CLIA**
 - Clinical Laboratory Improvement Amendments of 1988
- **CLFS**
 - Clinical Laboratory Fee Schedule
- **CPT**
 - Current Procedural Terminology

➢ Insurers typically have an annual limit for how many screens/tests can be performed

Painweek

<https://www.cmaa.org/Connect-and-Educate/Medical-Learning-Network/M/NM/Member/Article/Connect-and-Educate/1001.pdf>

29

UDM Billing

CPT Code AMA/CMS	UDS Presumptive/Qualitative Unlimited Drug Classes Includes Sample Validation	2018 Medicare Reimbursement Fee
80305	Immunoassay (Cup, Card, Cartridge, Dipstick) capable of being read by direct optical observation	\$13.46
80306	Immunoassay (Cup, Card, Cartridge, Dipstick) read by instrumented assisted direct optical observation	\$17.96
80307	Chemistry Analyzer (eg, EIA), Chromatography (eg, GC, HPLC), or Mass Spectrometry +/- Chromatography	\$71.83

Painweek

<https://www.cmaa.org/Connect-and-Educate/Medical-Learning-Network/M/NM/Member/Article/Connect-and-Educate/1001.pdf>

30

UDM Billing

UDM	CPT Code AMA/CMS	Description	2018 Medicare Reimbursement Fee
UDS	80307	Chemistry Analyzer (eg, EIA), Chromatography (eg, GC, HPLC), or Mass Spectrometry ± Chromatography	\$71.83
UDT	G0659	GC/LC with Mass Spectrometry in Physician's Office Lab (POL)	\$71.83



<https://www.cmaa.org/Outreach-and-Education/Medical-Learning/Network/M-NA-Medical-Clinical-Courtside-CE-18001.pdf>

31

UDM Billing

CPT Code AMA/CMS	UDT Definitive	2018 Medicare Reimbursement Fee
G0659	GC/LC with Mass Spectrometry in Physician's Office Lab (POL)	\$71.83
G0480	GC/LC with Mass Spectrometry with additional lab parameters (1 to 7 Drugs)	\$114.43
G0481	GC/LC with Mass Spectrometry with additional lab parameters (8 to 14 Drugs)	\$156.59
G0482	GC/LC with Mass Spectrometry with additional lab parameters (15 to 21 Drugs)	\$198.74
G0483	GC/LC with Mass Spectrometry with additional lab parameters (≥22 Drugs)	\$246.92



<https://www.cmaa.org/Outreach-and-Education/Medical-Learning/Network/M-NA-Medical-Clinical-Courtside-CE-18001.pdf>

32

UDM Positivity Rates

- Lab can provide your practice's rate of specific substances being positively identified in urine monitoring
- If 0% repeatedly for specific substances, one could possibly consider eliminating that urine monitoring
- However, drug utilization trends change and adapt over time



33

Profitable Pee

Maryland Medicaid program bans some drug testing due to costs



By Meredith Cahn • Contact Reporter
The Baltimore Sun

FEBRUARY 15, 2016, 3:30 PM

The greatest rivalries in criminal history are alive
National Law Enforcement



34

National Urine Drug Tests Overall Positive Results

Overall Positivity by 3-Digit Zip Code
Urine Drug Tests
January - December 2018

Percent Positive
■ 5.5% to 10%
■ 4.5% to 5.5%
■ 3.5% to 4.5%
■ 2.5% to 3.5%
■ 0.0% to 2.5%
□ Insufficient Data



35


National Urine Drug Tests PCP Positive Results

PCP Positivity by 3-Digit Zip Code
Urine Drug Tests
January - December 2018

Percent Positive
■ 0.1% to 0.5%
■ 0.02% to 0.1%
■ 0.01% to 0.03%
■ 0.001% to 0.01%
■ 0.0% to 0.001%
□ Insufficient Data




36




37



38



39



40

Cannabinoids


- THC metabolite: THC-COOH
 - Carboxylic acid group added to allow for kidney excretion
- CBD should not screen (+) for THC, however
 - High % of products contain other substances (eg, THC)
 - JAMA 2017 study: 26 of 84 (~30%) CBD extracts had accurate labels
 - Structures are very similar

CBD
CANNABIDIOL

THC
TETRAHYDROCANNABINOL

PAINweek

41



42

Patient Cases



Painweek

43

Patient Case #1

Prescribed oxycodone

- Conduct a pill count and complete history
- Pick UDS panel to evaluate adherence

Painweek

44

Patient Case #1

Prescribed oxycodone

- Conduct a pill count and complete history
- Pick UDS panel to evaluate adherence

URINE DRUG SCREENS				
SUBSTANCE	5 Panel	7 Panel	10 Panel	12 Panel
THC	X	X	X	X
Cocaine	X	X	X	X
Opiates	X	X	X	X
PCP	X	X	X	X
Amphetamines	X	X	X	X
Benzodiazepines		X	X	X
Barbiturates		X	X	X
Methadone			X	X
Propoxyphene			X	X
Quaaludes			X	X
Ecstasy				X
Oxycodone				X

Painweek

45

Patient Case #2

UDS 12 pain panel

- Lower than expected pill count
 - Diverted the rest?
 - Took a pill a few hours prior to UDS?
- Gradual (if any) opioid taper needed?

UDS 12 Pain Panel	Result
Marijuana (THC)	-
Cocaine	-
Opiates	-
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-
Methadone	-
Propoxyphene	-
Quaaludes	-
MDMA (Ecstasy)	-
OXYCODONE	+



46

Patient Case #2

UDS 12 pain panel

- Lower than expected pill count
 - Diverted the rest?
 - Took a pill a few hours prior to UDS?
- Gradual (if any) opioid taper needed?



UDT	Result
OXYCODONE	+
Oxymorphone	-
Noroxycodone	-

UDS 12 Pain Panel	Result
Marijuana (THC)	-
Cocaine	-
Opiates	-
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-
Methadone	-
Propoxyphene	-
Quaaludes	-
MDMA (Ecstasy)	-
OXYCODONE	+



47

Patient Case #3

- Verified current medication list (April)
 - Dextromethorphan, diphenhydramine, MVI, & lisinopril

UDS 7 Panel	Result
Marijuana (THC)	-
Cocaine	-
OPIATES	+
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-



48

Patient Case #3

- Verified current medication list (April)
 - Dextromethorphan, diphenhydramine, MVI, & lisinopril



UDT	Result
Morphine	-
6-MAM	-
Oxycodone	-
Oxymorphone	-
Noroxycodone	-
DEXTROMETHORPHAN	+
DIPHENHYDRAMINE	+

UDS 7 Panel	Result
Marijuana (THC)	-
Cocaine	-
OPIATES	+
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-



49

Structurally Speaking...



50

Dextromethorphan (DXM)


Dose (mg)	Effects
Low	Cough suppressant
Moderate	Pain management
100-200	Mild stimulation
201-400	Euphoria & hallucinations
540	Distorted vision & coordination
601-1500	Dissociative sedation

Drug Facts
 Active ingredient (in each 5 mL) Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide. Purpose Cough suppressant.




51

Chirality




Samuel-L-Jackson Samuel-D-Jackson



52

DXM

"Smartphone in Right Hand"



Dextromethorphan (DXM)

CN1CC2C(C1)C(C2)C

Methyl group

→

CYP-2D6

O-demethylation

Dextrorphan (DXO)
"D(+)" enantiomer of racemorphan


Right

↕

Levorphanol
"L(-)" enantiomer of racemorphan

MME: 11

Left



Clinical Pharmacology Online Database, 2016

53

Human Mind Maturing at Age 25...

Chain Drug Review

RETAIL NEWS SUPPLIER NEWS BUSINESS PHARMACY TECHNOLOGY OPINION LATEST ISSUES CDR BLOG VIDEOS SLIDESHOWS

West Virginia becomes 18th state to adopt age-18 sales law on cough medicine



https://www.painweek.com/news/industry-news/2016/04/20/wv-adopts-age-18-sales-law-on-cough-medicine/

54

Patient Case #4

- Patient currently not utilizing any medications

UDS 7 Panel	Result
Marijuana (THC)	-
Cocaine	-
OPIATES	+
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-

Painweek

55

Patient Case #4

- Patient currently not utilizing any medications
 - Aunt Doreen's Easter Bread
 - Poppy Seeds

UDT	Result
MORPHINE	+
6-MAM	-
Oxycodone	-
Oxymorphone	-
Noroxycodone	-

UDS 7 Panel	Result
Marijuana (THC)	-
Cocaine	-
OPIATES	+
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-

Painweek

56

“Poppy Seed Defense”

- Poppy seeds do not contain opium but are coated by it
- Concentration of coating varies, depending on
 - Amount of rainfall during the curing of the seed pods;
 - Washing of the seeds
- Bread, bagels, muffins, pastries, smoothies, fillings, etc
- Approximately 1 TSP poppy seeds may result in false-positive UDS for opiates
- Cake material: morphine urine concentrations can be in thousands of ng/mL²
- Bagels: morphine urine concentrations can be in hundreds of ng/mL³

Painweek

1. https://pubmed.ncbi.nlm.nih.gov/20199688
 2. Thomas M, Calkins G, and Calhoun W. Urinary Concentrations of Morphine and Codeine After Consumption of Poppy Seeds. J Anal Toxicol. 2003;27
 3. Bostic TP, Moore G. The Determination of Morphine in Urine and Oral Fluid Following Ingestion of Poppy Seeds. J Anal Toxicol. 2004;28

57

“Poppy Seed Defense”

2015 Mayo Clinic Case Study

Patient reported drinking tea prepared from 1 to 2 POUNDS of poppy seeds daily (OWS upon discontinuation). The patient’s LCMS results included:

- Morphine 37,600 ng/mL
- Codeine 2,580 ng/mL
- Hydromorphone 1,430 ng/mL

- Thebaine: Short T1/2 (similar to 6-MAM)
- Codeine & Morphine produce <11% hydrocodone & <2.5% hydromorphone
 - >These 2 metabolites have not been reported with poppy seed use
 - >1,430/37,600 = 3.8% (In other words, >2.5%)



Patenaik A, et al. Interpreting Urine Drug Screen Results in the Context of Poppy Seed Use. Mayo Clinic Proceedings. 2015;92(12):1712-1715.

58

Patient Case #5

- Broken arm, patient not happy about needing to conduct a UDS due to recent UTI
- You ask how long ago the UTI was, and patient says it still hurts

Which of the following antibiotics could produce a false positive UDS for opiates?

- a) Azithromycin
- b) Cefdinir
- c) Cephalexin
- d) Ciprofloxacin



59

Patient Case #5

- Broken arm, patient not happy about needing to conduct a UDS due to recent UTI
- You ask how long ago the UTI was, and patient says it still hurts

Which of the following antibiotics could produce a false positive UDS for opiates?

- a) Azithromycin
- b) Cefdinir
- c) Cephalexin
- d) CIPROFLOXACIN*



60

Patient Case #6

- Patient currently not utilizing any medications

UDS 7 Panel	Result
MARIJUANA (THC)	+
Cocaine	-
Opiates	-
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-

Painweek

61

Patient Case #6

- Patient currently not utilizing any medications
- Ibuprofen 200 mg PRN
- Didn't remember buying at The Dollar Store
- Recently taken while gardening

UDT	Result
THC	-
Amphetamine	-
Morphine	-
Oxycodone	-

UDS 7 Panel	Result
MARIJUANA (THC)	+
Cocaine	-
Opiates	-
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-

Painweek

62

Patient Case #7

- Patient attempted to falsify the UDS by ingesting goldenseal
- Goldenseal is an herb in the buttercup family with a large yellow root, native in southeastern Canada and the northeastern United States.

UDT	Result
THC	+
Amphetamine	-
Morphine	-
Oxycodone	-

UDS 7 Panel	Result
MARIJUANA (THC)	+
Cocaine	-
Opiates	-
PCP	-
Amphetamines	-
Benzodiazepines	-
Barbiturates	-

Painweek

63

Goldenseal: Myth, Legend, Fact, Fiction?

- John Uri Lloyd, a pharmacist, wrote the fiction novel "Stringtown on the Pike" in 1890, involving a murder with a victim having strychnine in the stomach.
- The murder victim's morning habit of taking bitters (an ingredient of which was goldenseal), was the cause of the false positive for the deadly poison.
- Two recent scientific studies have showed no effect of goldenseal on urine drug tests. Subjects who drank large amounts of water had the same urine drug levels as subjects who took goldenseal pills with the water.



64

Goldenseal: Myth, Legend, Fact, Fiction?

I have to take a drug test on MOnday and last time I smoked was Wed night. U usually smoke a couple times a week- maybe a joint here and there.

the test is called 35105N SAP 5-50 w/nit

So, my fiance told me golden seal worked for him before. He told me to drink tons of water today and tomorrow and then just regularly on Sunday so I'm not overhydrated. Will this work??

If it helps, I'm 6'1, 165lbs and exercises regularly (2 to 3times per week) and am female.

Thanks a ton if anyone can help me!

lustywrench, May 21, 2004



65

Goldenseal: Myth, Legend, Fact, Fiction?

so i passed

So, I passed the test, cause I start my job tomorrow...thanks for the advice everyone

FYI- I did take golden seal and drank over 100 ounces water fri and sat. each and ate meat. Sunday I didn't overdrink water but started taking aspirin regularly. Then Mon. before test I took aspirin 4 hours before test and drank a powerade an hour before. it worked so I don't know which part was helpful, but frankly at this point I don't care cause I got JOB now!

Thanks

lustywrench, May 25, 2004



66

Patient Case #8

- Confirmed current medication list
 - Acetaminophen/codeine #4 Q 4 hours PRN
 - Ibuprofen 200mg Q 6 hours PRN
- Pharmacogenomics (CYP450-2D6)?
- “Overuse” due to sub-potent opioid?
- Illicit use of other opioids?




UDT	Result
Metadone	-
EDDP	-
MORPHINE	+
6-MAM	-
HYDROMORPHONE	+
HYDROCODONE	+
DIHYDROCODEINE	+
Oxycodone	-
Oxymorphone	-
Noroxycodone	-

Painweek


67

Opioid Metabolism
Active Metabolites



Painweek

68



Painweek

69

Patient Dismissal



Painweek

70

Confirmed Drug Seeking or Diversion

- Reference the patient and provider agreement/contract
- Treatment can continue with noncontrolled substance therapies
- Refer to a substance use disorder specialist/program, or an entity that can connect
- Contact law enforcement if concern for the safety of the patient or others exists
- Respect for all those directly or indirectly involved in the specific patient case should be upheld, while also ensuring both a procession within federal/state laws and an appropriate level of patient care

Painweek

71

Avoid Patient Abandonment

1. DOCUMENT everything thoroughly
2. Patient signs termination letter in person with a witness (or certified mail return receipt)
 - Provide at least a 30-day notice in the letter and offer to continue medical treatment for that time period
 - Stress the importance of continued medical care and potential consequences of not getting that care
 - Refer the patient to a number of professionals whom may accept the patient for care
 - Offer to promptly transfer medical records to the new provider once established

Painweek

Palmer D, et al. Avoid Medical Abandonment: Chronic Opioid Analgesic Therapy. Case Report. Pain Medicine Volume 10, Issue 4, 4 May 2009. Pages 770-781

72

https://www.deadiversion.usdoj.gov/tips_online.htm

1-877-RX-Abuse
(1-877-792-2873)

73

WVU Medicine Center for Integrative Pain Management

- ALL patients have a UDT (GCMS)
- Results available within 24 hours
- Results provided via secure online portal website
 - Preset notification settings (timing, specific results, etc)
- Violators receive a certified letter
- Conversation....

Painweek

<https://www.wvu.edu/~medcenter/hospital/clinics/specialty-clinics/pain-management-center>

74

To Infinity, and Beyond...

Painweek

75

Audience Question #1

A urine drug screening (UDS) is confirmatory and quantitative, whereas a urine drug testing (UDT) is presumptive and qualitative.

- a) True
- b) False



76

Audience Question #1 (ANSWER)

A urine drug screening (UDS) is confirmatory and quantitative, whereas a urine drug testing (UDT) is presumptive and qualitative.

- a) True
- b) FALSE [CORRECT ANSWER]



77

Audience Question #2

A 35yo patient with chronic lower back pain, well managed with PT, CBT, and OTC naproxen 220 mg 3 times daily. Patient screens opiate +, yet denies opioid utilization. Patient mentions having utilized a nighttime over-the-counter medication to help sleep with a cold recently. Which of the following OTC medications could cause of a false opiate + UDS result?

- a) Only dextromethorphan
- b) Only diphenhydramine
- c) Both dextromethorphan and diphenhydramine
- d) Neither dextromethorphan nor diphenhydramine



78

Audience Question #2 (ANSWER)

A 35yo patient with chronic lower back pain, well managed with PT, CBT, and OTC naproxen 220 mg 3 times daily. Patient screens opiate +, yet denies opioid utilization. Patient mentions having utilized a nighttime over-the-counter medication to help sleep with a cold recently. Which of the following OTC medications could cause of a false opiate + UDS result?

- a) Only dextromethorphan
- b) Only diphenhydramine
- c) BOTH DEXTROMETHORPHAN AND DIPHENHYDRAMINE
[CORRECT ANSWER]
- d) Neither dextromethorphan nor diphenhydramine



79

Audience Question #3

Which of the following medications can possibly produce a false positive urine drug screenings for marijuana (THC)?

- a) Ibuprofen
- b) Omeprazole
- c) Promethazine
- d) All of the above



80

Audience Question #3 (ANSWER)

Which of the following medications can possibly produce a false positive urine drug screenings for marijuana (THC)?

- a) Ibuprofen
- b) Omeprazole
- c) Promethazine
- d) ALL OF THE ABOVE [CORRECT ANSWER]



81

**PAINWeek
National Conference
September 7-11 in Las Vegas**

Special Offer!
Save \$100 off registration:
Go to painweek.org and use code PWP21.

This discount does not apply to industry registrations, or those already registered for PAINWeek 2021, and cannot be combined with other discounts.